

Systems Of Equations,

Definition 1.2: System of Linear Equations

A **system of linear equations** is a list of equations,

$$\begin{aligned}a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n &= b_1 \\a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n &= b_2 \\&\vdots \\a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n &= b_m\end{aligned}$$

where a_{ij} and b_j are real numbers. The above is a system of m equations in the n variables, x_1, x_2, \dots, x_n . Written more simply in terms of summation notation, the above can be written in the form

$$\sum_{j=1}^n a_{ij}x_j = b_i, \quad i = 1, 2, 3, \dots, m$$

Definition 1.3: Homogeneous System of Equations

A system of equations is called **homogeneous** if each equation in the system is equal to 0. A homogeneous system has the form

$$\begin{aligned}a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n &= 0 \\a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n &= 0 \\&\vdots \\a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n &= 0\end{aligned}$$

where a_{ij} are scalars and x_i are variables.

Definition 1.4: Consistent and Inconsistent Systems

A system of linear equations is called **consistent** if there exists at least one solution. It is called **inconsistent** if there is no solution.

Elementary Operations

Definition 1.6: Elementary Operations

Elementary operations are those operations consisting of the following.

1. Interchange the order in which the equations are listed.
2. Multiply any equation by a nonzero number.
3. Replace any equation with itself added to a multiple of another equation.

Theorem 1.8: Elementary Operations and Solutions

Suppose you have a system of two linear equations

$$\begin{aligned}E_1 &= b_1 \\E_2 &= b_2\end{aligned}\tag{1.1}$$

Then the following systems have the same solution set as 1.1:

1.

$$\begin{aligned}E_2 &= b_2 \\E_1 &= b_1\end{aligned}\tag{1.2}$$

2.

$$\begin{aligned}E_1 &= b_1 \\kE_2 &= kb_2\end{aligned}\tag{1.3}$$

for any scalar k , provided $k \neq 0$.

3.

$$\begin{aligned}E_1 &= b_1 \\E_2 + kE_1 &= b_2 + kb_1\end{aligned}\tag{1.4}$$

for any scalar k (including $k = 0$).

Example 1.9: Solving a System of Equations with Elementary Operations

Find the solutions to the system,

$$\begin{aligned}x + 3y + 6z &= 25 \\2x + 7y + 14z &= 58 \\2y + 5z &= 19\end{aligned}\tag{1.5}$$

Solution. We can relate this system to Theorem 1.8 above. In this case, we have

$$\begin{aligned}E_1 &= x + 3y + 6z, & b_1 &= 25 \\E_2 &= 2x + 7y + 14z, & b_2 &= 58 \\E_3 &= 2y + 5z, & b_3 &= 19\end{aligned}$$

$$\begin{aligned}x &= 1 \\y &= 2 \\z &= 3\end{aligned}$$

Gaussian Elimination

Definition 1.10: Augmented Matrix of a Linear System

For a linear system of the form

$$\begin{aligned} a_{11}x_1 + \cdots + a_{1n}x_n &= b_1 \\ &\vdots \\ a_{m1}x_1 + \cdots + a_{mn}x_n &= b_m \end{aligned}$$

where the x_i are variables and the a_{ij} and b_i are constants, the augmented matrix of this system is given by

$$\left[\begin{array}{ccc|c} a_{11} & \cdots & a_{1n} & b_1 \\ \vdots & & \vdots & \vdots \\ a_{m1} & \cdots & a_{mn} & b_m \end{array} \right]$$

Definition 1.11: Elementary Row Operations

The **elementary row operations** (also known as **row operations**) consist of the following

1. Switch two rows.
2. Multiply a row by a nonzero number.
3. Replace a row by any multiple of another row added to it.

For example, consider the linear system in Example 1.9

$$\begin{aligned} x + 3y + 6z &= 25 \\ 2x + 7y + 14z &= 58 \\ 2y + 5z &= 19 \end{aligned}$$

This system can be written as an augmented matrix, as follows

$$\left[\begin{array}{ccc|c} 1 & 3 & 6 & 25 \\ 2 & 7 & 14 & 58 \\ 0 & 2 & 5 & 19 \end{array} \right]$$

$$\left[\begin{array}{ccc|c} 1 & 3 & 6 & 25 \\ 0 & 1 & 2 & 8 \\ 0 & 2 & 5 & 19 \end{array} \right] \quad \left[\begin{array}{ccc|c} 1 & 3 & 6 & 25 \\ 0 & 1 & 2 & 8 \\ 0 & 0 & 1 & 3 \end{array} \right]$$

This augmented matrix corresponds to the system

$$\begin{aligned} x + 3y + 6z &= 25 \\ y + 2z &= 8 \\ z &= 3 \end{aligned}$$

which is the same as 1.7. By back substitution you obtain the solution $x = 1, y = 2$, and $z = 3$.

Definition 1.12: Row-Echelon Form

An augmented matrix is in **row-echelon form** if

1. All nonzero rows are above any rows of zeros.
2. Each leading entry of a row is in a column to the right of the leading entries of any row above it.
3. Each leading entry of a row is equal to 1.

Definition 1.13: Reduced Row-Echelon Form

An augmented matrix is in **reduced row-echelon form** if

1. All nonzero rows are above any rows of zeros.
2. Each leading entry of a row is in a column to the right of the leading entries of any rows above it.
3. Each leading entry of a row is equal to 1.
4. All entries in a column above and below a leading entry are zero.

Example 1.14: Not in Row-Echelon Form

The following augmented matrices are not in row-echelon form (and therefore also not in reduced row-echelon form).

$$\left[\begin{array}{ccc|c} 0 & 0 & 0 & 0 \\ 1 & 2 & 3 & 3 \\ 0 & 1 & 0 & 2 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{array} \right], \left[\begin{array}{cc|c} 1 & 2 & 3 \\ 2 & 4 & -6 \\ 4 & 0 & 7 \end{array} \right], \left[\begin{array}{ccc|c} 0 & 2 & 3 & 3 \\ 1 & 5 & 0 & 2 \\ 7 & 5 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{array} \right]$$

Example 1.15: Matrices in Row-Echelon Form

The following augmented matrices are in row-echelon form, but not in reduced row-echelon form.

$$\left[\begin{array}{ccccc|c} 1 & 0 & 6 & 5 & 8 & 2 \\ 0 & 0 & 1 & 2 & 7 & 3 \\ 0 & 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{array} \right], \left[\begin{array}{ccc|c} 1 & 3 & 5 & 4 \\ 0 & 1 & 0 & 7 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{array} \right], \left[\begin{array}{ccc|c} 1 & 0 & 6 & 0 \\ 0 & 1 & 4 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

Example 1.16: Matrices in Reduced Row-Echelon Form

The following augmented matrices are in reduced row-echelon form.

$$\left[\begin{array}{ccccc|c} 1 & 0 & 0 & 5 & 0 & 0 \\ 0 & 0 & 1 & 2 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{array} \right], \left[\begin{array}{ccc|c} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{array} \right], \left[\begin{array}{ccc|c} 1 & 0 & 0 & 4 \\ 0 & 1 & 0 & 3 \\ 0 & 0 & 1 & 2 \end{array} \right]$$

Definition 1.17: Pivot Position and Pivot Column

A **pivot position** in a matrix is the location of a leading entry in the row-echelon form of a matrix. A **pivot column** is a column that contains a pivot position.

Example 1.18: Pivot Position

Let

$$A = \left[\begin{array}{ccc|c} 1 & 2 & 3 & 4 \\ 3 & 2 & 1 & 6 \\ 4 & 4 & 4 & 10 \end{array} \right]$$

Where are the pivot positions and pivot columns of the augmented matrix A ?

Solution. The row-echelon form of this matrix is

$$\left[\begin{array}{ccc|c} 1 & 2 & 3 & 4 \\ 0 & 1 & 2 & \frac{3}{2} \\ 0 & 0 & 0 & 0 \end{array} \right]$$

$$\left[\begin{array}{ccc|c} \boxed{1} & 2 & 3 & 4 \\ 3 & \boxed{2} & 1 & 6 \\ 4 & 4 & 4 & 10 \end{array} \right]$$

Thus the pivot columns in the matrix are the first two columns.

Algorithm 1.19: Reduced Row-Echelon Form Algorithm

This algorithm provides a method for using row operations to take a matrix to its reduced row-echelon form. We begin with the matrix in its original form.

1. Starting from the left, find the first nonzero column. This is the first pivot column, and the position at the top of this column is the first pivot position. Switch rows if necessary to place a nonzero number in the first pivot position.
2. Use row operations to make the entries below the first pivot position (in the first pivot column) equal to zero.
3. Ignoring the row containing the first pivot position, repeat steps 1 and 2 with the remaining rows. Repeat the process until there are no more rows to modify.
4. Divide each nonzero row by the value of the leading entry, so that the leading entry becomes 1. The matrix will then be in row-echelon form.

The following step will carry the matrix from row-echelon form to reduced row-echelon form.

5. Moving from right to left, use row operations to create zeros in the entries of the pivot columns which are above the pivot positions. The result will be a matrix in reduced row-echelon form.

Example 1.20: Finding Row-Echelon Form and Reduced Row-Echelon Form of a Matrix

Let

$$A = \begin{bmatrix} 0 & -5 & -4 \\ 1 & 4 & 3 \\ 5 & 10 & 7 \end{bmatrix}$$

Find the row-echelon form of A . Then complete the process until A is in reduced row-echelon form.

Solution. In working through this example, we will use the steps outlined in Algorithm 1.19.

1. The first pivot column is the first column of the matrix, as this is the first nonzero column from the left. Hence the first pivot position is the one in the first row and first column. Switch the first two rows to obtain a nonzero entry in the first pivot position, outlined in a box below.

$$\begin{bmatrix} \boxed{1} & 4 & 3 \\ 0 & -5 & -4 \\ 5 & 10 & 7 \end{bmatrix}$$

2. Step two involves creating zeros in the entries below the first pivot position. The first entry of the second row is already a zero. All we need to do is subtract 5 times the first row from the third row. The resulting matrix is

$$\begin{bmatrix} 1 & 4 & 3 \\ 0 & -5 & -4 \\ 0 & 10 & 8 \end{bmatrix}$$

3. Now ignore the top row. Apply steps 1 and 2 to the smaller matrix

$$\begin{bmatrix} -5 & -4 \\ 10 & 8 \end{bmatrix}$$

In this matrix, the first column is a pivot column, and -5 is in the first pivot position. Therefore, we need to create a zero below it. To do this, add 2 times the first row (of this matrix) to the second. The resulting matrix is

$$\begin{bmatrix} -5 & -4 \\ 0 & 0 \end{bmatrix}$$

Our original matrix now looks like

$$\begin{bmatrix} 1 & 4 & 3 \\ 0 & -5 & -4 \\ 0 & 0 & 0 \end{bmatrix}$$

We can see that there are no more rows to modify.

4. Now, we need to create leading 1s in each row. The first row already has a leading 1 so no work is needed here. Divide the second row by -5 to create a leading 1. The resulting matrix is

$$\begin{bmatrix} 1 & 4 & 3 \\ 0 & 1 & \frac{4}{5} \\ 0 & 0 & 0 \end{bmatrix}$$

This matrix is now in row-echelon form.

5. Now create zeros in the entries above pivot positions in each column, in order to carry this matrix all the way to reduced row-echelon form. Notice that there is no pivot position in the third column so we do not need to create any zeros in this column! The column in which we need to create zeros is the second. To do so, subtract 4 times the second row from the first row. The resulting matrix is

$$\begin{bmatrix} 1 & 0 & -\frac{1}{5} \\ 0 & 1 & \frac{4}{5} \\ 0 & 0 & 0 \end{bmatrix}$$

This matrix is now in reduced row-echelon form.

Example 1.21: Finding the Solution to a System

Give the complete solution to the following system of equations

$$\begin{aligned} 2x + 4y - 3z &= -1 \\ 5x + 10y - 7z &= -2 \\ 3x + 6y + 5z &= 9 \end{aligned}$$

Solution. The augmented matrix for this system is

$$\left[\begin{array}{ccc|c} 2 & 4 & -3 & -1 \\ 5 & 10 & -7 & -2 \\ 3 & 6 & 5 & 9 \end{array} \right]$$

$$\left[\begin{array}{ccc|c} 2 & 4 & -3 & -1 \\ 0 & 0 & 1 & 1 \\ 3 & 6 & 5 & 9 \end{array} \right] \quad \left[\begin{array}{ccc|c} 2 & 4 & -3 & -1 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 1 & 21 \end{array} \right] \quad \left[\begin{array}{ccc|c} 2 & 4 & -3 & -1 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 20 \end{array} \right]$$

$$0x + 0y + 0z = 20$$

There is no solution to this equation because for all x, y, z , the left side will equal 0 and $0 \neq 20$. This shows there is no solution to the given system of equations. In other words, this system is inconsistent. ♠

Example 1.22: An Infinite Set of Solutions

Give the complete solution to the system of equations

$$\begin{aligned} 3x - y - 5z &= 9 \\ y - 10z &= 0 \\ -2x + y &= -6 \end{aligned} \tag{1.8}$$

Solution. The augmented matrix of this system is

$$\left[\begin{array}{ccc|c} 3 & -1 & -5 & 9 \\ 0 & 1 & -10 & 0 \\ -2 & 1 & 0 & -6 \end{array} \right]$$

$$\left[\begin{array}{ccc|c} 3 & -1 & -5 & 9 \\ 0 & 1 & -10 & 0 \\ 0 & 1 & -10 & 0 \end{array} \right] \quad \left[\begin{array}{ccc|c} 3 & -1 & -5 & 9 \\ 0 & 1 & -10 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right] \quad \left[\begin{array}{ccc|c} 1 & -\frac{1}{3} & -\frac{5}{3} & 3 \\ 0 & 1 & -10 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

This matrix is now in row-echelon form.

$$\left[\begin{array}{ccc|c} 1 & 0 & -5 & 3 \\ 0 & 1 & -10 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

This is in reduced row-echelon form, which you should verify using Definition 1.13. The equations corresponding to this reduced row-echelon form are

$$\begin{aligned} x - 5z &= 3 \\ y - 10z &= 0 \end{aligned}$$

or

$$\begin{aligned} x &= 3 + 5z \\ y &= 10z \end{aligned}$$

let $z = t$, where we can choose t to be any number. In this context t is called a **parameter**. Therefore, the solution set of this system is

$$\begin{aligned} x &= 3 + 5t \\ y &= 10t \\ z &= t \end{aligned}$$

Example 1.23: A Two Parameter Set of Solutions

Find the solution to the system

$$\begin{aligned} x + 2y - z + w &= 3 \\ x + y - z + w &= 1 \\ x + 3y - z + w &= 5 \end{aligned}$$

Solution. The augmented matrix is

$$\left[\begin{array}{cccc|c} 1 & 2 & -1 & 1 & 3 \\ 1 & 1 & -1 & 1 & 1 \\ 1 & 3 & -1 & 1 & 5 \end{array} \right]$$

$$\left[\begin{array}{cccc|c} 1 & 2 & -1 & 1 & 3 \\ 0 & -1 & 0 & 0 & -2 \\ 0 & 1 & 0 & 0 & 2 \end{array} \right] \quad \left[\begin{array}{cccc|c} 1 & 2 & -1 & 1 & 3 \\ 0 & 1 & 0 & 0 & 2 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right]$$

$$\begin{aligned} x &= -1 + s - t \\ y &= 2 \\ z &= s \\ w &= t \end{aligned}$$

It is customary to write this solution in the form

$$\begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix} = \begin{bmatrix} -1 + s - t \\ 2 \\ s \\ t \end{bmatrix}$$

Definition 1.33: Rank of a Matrix

Let A be a matrix and consider any row-echelon form of A . Then, the number r of leading entries of A does not depend on the row-echelon form you choose, and is called the **rank** of A . We denote it by $\text{rank}(A)$.

Similarly, we could count the number of pivot positions (or pivot columns) to determine the rank of A .

Example 1.34: Finding the Rank of a Matrix

Consider the matrix

$$\begin{bmatrix} 1 & 2 & 3 \\ 1 & 5 & 9 \\ 2 & 4 & 6 \end{bmatrix}$$

What is its rank?

Solution. First, we need to find the reduced row-echelon form of A . Through the usual algorithm, we find that this is

$$\begin{bmatrix} \boxed{1} & 0 & -1 \\ 0 & \boxed{1} & 2 \\ 0 & 0 & 0 \end{bmatrix}$$

Here we have two leading entries, or two pivot positions, shown above in boxes. The rank of A is $r = 2$.

Theorem 1.35: Rank and Solutions to a Homogeneous System

Let A be the $m \times n$ coefficient matrix corresponding to a homogeneous system of equations, and suppose A has rank r . Then, the solution to the corresponding system has $n - r$ parameters.

Theorem 1.36: Rank and Solutions to a Consistent System of Equations

Let A be the $m \times (n + 1)$ augmented matrix corresponding to a consistent system of equations in n variables, and suppose A has rank r . Then

1. the system has a unique solution if $r = n$
2. the system has infinitely many solutions if $r < n$